



## Recommended Immunizations for Children from Birth Through 6 Years Old

Birth	2 Months	4 Months	6 Months	9 Months	12 Months	15 Months	18 Months	2-3 Years	4-6 Years
Hep B	Hep B		Hep B						
	Rota	Rota	Rota						
	DTaP	DTaP	DTaP				DTaP		DTaP
	Hib	Hib	Hib			Hib			
	PCV	PCV	PCV			PCV <sup>1</sup>			
	IPV	IPV	IPV						IPV
						MMR			MMR
						Varicella			Varicella
						Hep A <sup>3</sup>			
			Influenza (yearly) <sup>2</sup>						

### Footnotes:

If your child misses a shot, you don't need to start over, just go back to your child's doctor for the next shot. The doctor will keep your child up-to-date on vaccinations. Talk with your doctor if you have questions.

<sup>1</sup>Children 2 years old and older with certain medical conditions may need a dose of pneumococcal vaccine (PPSV) and meningococcal vaccine (MCV4). See vaccine-specific recommendations at <http://www.cdc.gov/vaccines/pubs/ACIP-list.htm>.

<sup>2</sup>Two doses given at least four weeks apart are recommended for children aged 6 months through 8 years of age who are getting a flu vaccine for the first time.

<sup>3</sup>Two doses of HepA vaccine are needed for lasting protection. The first dose of HepA vaccine should be given between 12 months and 23 months of age. The second dose should be given 6 to 18 months later. HepA vaccination may be given to any child 12 months and older to protect against HepA. Children and adolescents who did not receive the HepA vaccine and are at high-risk, should be vaccinated against HepA.



American Academy of Pediatrics  
DEDICATED TO THE HEALTH OF ALL CHILDREN™



For more information, call toll free 1-800-CDC-INFO (432-695-6300) or visit <http://www.cdc.gov/vaccines/teens>

**PEDCARE**  
Children's Clinic

Address : 3001 W Illinois Ave Ste # 1A, Midland, TX 79701